

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 June 2003 (19.06.2003)

PCT

(10) International Publication Number
WO 03/049752 A2

(51) International Patent Classification⁷: **A61K 35/74**,
39/04, A61P 35/00, 37/00, 31/06

(21) International Application Number: **PCT/IB02/05760**

(22) International Filing Date:
11 December 2002 (11.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01403195.9 11 December 2001 (11.12.2001) EP

(71) Applicants (*for all designated States except US*): **INSTITUT PASTEUR** [FR/FR]; 28 rue du Docteur Roux, F-75724 Cedex 15 Paris (FR). **INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE** [FR/FR]; 101 rue de Tolbiac, F-75654 Cedex 13 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **NAHORI, Marie-Anne** [FR/FR]; 147 avenue Henri Ginoux, F-92120 Montrouge (FR). **LAGRANDERIE, Micheline** [FR/FR]; 84 rue Charles Laffitte, F-92200 Neuilly-sur-Seine (FR). **MARCHAL, Gilles** [FR/FR]; 4 rue Francisco Ferrer, F-94200 Ivry-sur-Seine (FR). **VARGAFTIG, Bernardo, Boris** [FR/FR]; 5 Villa Poirier, F-75015 Paris (FR). **LEFORT, Jean** [FR/FR]; 2 rue des Cailloux, F-92110 Clichy-sur-Seine (FR). **ROMAIN, Félix** [FR/FR]; 49

bis rue Dreyfus, F-91640 Fontenay les Briis (FR). **HEKIMIAN, Georges** [FR/FR]; 91 rue Jean Mermoz, F-92380 Garches (FR). **PELTRE, Philippe** [FR/FR]; 100 Boulevard Georges Seurat, F-92200 Neuilly-sur-Seine (FR).

(74) Agent: **CABINET ORES**; 36 rue de St Petersburg, F-75008 Paris (FR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GRAM POSITIVE BACTERIA PREPARATIONS FOR THE TREATMENT OF DISEASES COMPRISING AN IMMUNE DYSREGULATION

(57) Abstract: Compositions comprising components prepared from Gram positive bacteria such as Gram positive facultative intracellular bacteria, for example mycobacteria, including extended freeze-dried killed Gram positive bacteria, their method of preparation and their use in humans and animals, for the prevention and/or the treatment of disorders comprising an immune dysregulation such as cancer, autoimmune diseases, allergy and tuberculosis.

BEST AVAILABLE COPY

WO 03/049752 A2

GRAM POSITIVE BACTERIA PREPARATIONS FOR THE TREATMENT OF DISEASES COMPRISING AN IMMUNE DYSREGULATION

The present invention relates to compositions comprising components prepared from Gram positive bacteria such as Gram positive facultative intracellular bacteria, for example mycobacteria, for the treatment of disorders comprising an immune dysregulation, in humans and animals. The invention also relates to the preparation of said components and compositions.

Diseases comprising an immune dysregulation, such as a Th1-Th2 imbalance, include different cancers, autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, Crohn disease and diabetes mellitus, and allergic disorders such as asthma, allergic rhinitis, conjunctivitis and atopic dermatitis.

For instance:

Allergic asthma is a common disease, which involves airway allergen-induced inflammation; the inflammatory leukocytes, eosinophils and sometimes neutrophils, are recruited to airways as a consequence of the presence and activation of T lymphocytes that react with the inhaled allergens in the respiratory tract. CD4⁺ T lymphocytes play a major role in initiating allergic airways inflammation via the production of Th2-type cytokines, which trigger the recruitment of eosinophils to the airways and possibly their subsequent activation. It has been proposed that an imbalance between Th2 and Th1 effectors drives the pathogenesis of asthma.

Therefore, it has been proposed to stimulate a Th1 immune response in the lungs, for instance by nebulization of IFN- γ or by administration of a mycobacterial vaccine; such a stimulation seems to inhibit the development of secondary allergic processes in mice.

More specifically, extrinsic asthma or atopic asthma (e.g. occupational and drug-induced) is associated with the enhancement of a Th2 type immune response with the production of specific immunoglobulin E (IgE), positive skin tests to common aeroallergens and/or atopic symptoms. The airflow obstruction in extrinsic asthma involves a non-specific bronchial hyperresponsiveness (BHR) caused by inflammation of the airways. This inflammation is mediated by chemicals released by a variety of inflammatory cells including mast cells, eosinophils and lymphocytes. The

actions of these mediators result in vascular permeability, mucus secretion and bronchial smooth muscle constriction. In atopic asthma, the immune response producing airway inflammation is brought about by Th2 class of T cells which secrete IL-4 and IL-5. Intrinsic or cryptogenic asthma is reported to develop after upper respiratory tract infections, but can arise *de novo* in middle-aged or older people, in whom it is more difficult to treat than extrinsic asthma.

Therefore allergic disorders are mediated by T lymphocytes secreting T helper 2 (Th2) cytokines, IL-4, IL-5 and IL-13, resulting in high levels of serum IgE and recruitment of eosinophils. However, Th1 cells may also participate in these pathologies; for instance, Th1 cells have been identified in chronic atopic skin lesions. Such a Th1-Th2 imbalance has also been found in other disorders such as some cancers, especially bladder cancer or in autoimmune diseases.

Epidemiological studies, for instance those performed by Shirakawa and coll. (Science, 1997, 275, 3), demonstrated an inverse association between tuberculin response and atopic disorders. Such results indicate that previous immunization against certain living Gram positive bacteria such as Gram positive facultative intracellular bacteria, for example mycobacteria such as *M. tuberculosis*, *M. bovis* or BCG, can protect against atopic diseases.

One of the treatment strategies is to down regulate the Th2 component by inducing a T helper 1 (Th1) response to the relevant allergen or antigen, because Th1 and Th2 cytokines are thought to be mutually antagonistic.

Experimental results from different groups have shown that the immunisation of adult mice with intracellular bacteria including mycobacteria and listeria, known to induce a strong Th1 immune response, may counterbalance the allergen or antigen induced Th2 response; in allergy, for instance, it may reduce eosinophilia and the associated BHR.

It is the reason why certain mycobacteria have been proposed for treating allergic disorders and more specifically asthma.

In this respect, different kind of compositions including mycobacteria have been tested:

- live BCG vaccines have been considered as having a potential effect in experimental asthma or equivalent, specially when given intranasally (KJ. Erb and coll., J. Exp. Med, 1998, **187**, 561-569; MA. Nahori and coll., Vaccine, 2001, **19**, 1484-1495). However, despite their utility to prevent tuberculosis (JB Milstein and coll., WHO Bull. OMS, 1990, **68**, 93-108), live BCG vaccines exhibit several drawbacks; first, they cannot be given to immuno-compromised subjects due to their residual virulence; second, there is a local reaction to intradermal BCG vaccination which is proportional to the total bacterial mass and can lead to local ulceration, or in the case of accidental subcutaneous injection, to more severe reactions (abscess). Therefore, such a vaccine, especially if it was given by the intranasal or aerosol route, would not be adapted for repeated administration over a significant period for the immunotherapy of allergic disorders of the respiratory tract, or other disorders involving a Th1-Th2 imbalance. Administration intranasally or by aerosol could lead to inadmissible adverse effects of the live BCG vaccine in the lungs;

- heat-killed preparations of mycobacteria:

. heat-killed preparations of BCG or of *Mycobacterium tuberculosis* do not have systematic protective properties when employed in vaccines (R. Janssen and coll., *Immunol.*, 2001, **102**, 4, 441-449; MA Skinner and coll., *Immunol.*, 2001, **102**, 2, 225-233; GA Rook and coll., Novartis Found Symposium, 1998, **217**, 73-87 and 87-98). Moreover, said preparations induce a delayed type hypersensitivity to BCG purified protein derivatives (PPD) that interferes with the diagnosis of tuberculosis. However, it must be noted that some authors have considered that pre-immunization in the newborn period with heat-killed *M. bovis* alone or in addition to *M. vaccae* may potentially be helpful in down-regulating an IgE response (F. Tukenmez and coll., *Pediatr. Allergy Immunol.*, 1999, **10**, 2, 107-111; F. Tukenmez and coll., J. Asthma, 2000, **37**, 4, 329-334; Nahori and coll., precited);

. heat-killed preparations of *Mycobacterium vaccae*, (CC Wang and coll., *Immunol.* 1998, **93**, 3, 307-313; WO 00/74715) have clinical application in the immunotherapy of allergy. Some results demonstrate the potency of heat-killed mycobacteria as Th1 adjuvants and show a potential application for recombinant mycobacteria in antigen-specific immune modulation (R. Janssen and coll., *Immunol.*, 2001,

102, 4, 441-449; MA Skinner and coll., *Immunol.*, 2001, 102, 2, 225-233; CC Wang and coll., *Immunol.* 1998, 93, 3, 307-313; WO 00/74715). Even though, said preparations are active in immunotherapy, they present several drawbacks, shared with the heat-killed preparations of BCG or of *Mycobacterium tuberculosis*, which also
5 preclude their use in immunotherapy.

Indeed, all the heat-killed preparations of mycobacteria:

- induce the production of TNF- α by alveolar macrophages, which behaves as a toxic and necrotizing substance, this precluding the use of such heat-killed preparations in immunotherapy, and
- 10 - induce local side effects (PM Shirtcliffe and coll., *Am. J. Respir. Crit. Care Med.*, 2001, 163, 1410-1414), such as itch blistering, induration and necrosis with persisting scares observed in human clinical trials.

Therefore, there is a need for efficient compositions derived from Gram positive bacteria such as Gram positive facultative intracellular bacteria, for
15 example mycobacteria, for the treatment of diseases comprising an immune dysregulation such as a Th1-Th2 imbalance, for example allergic disorders, said compositions being well-tolerated and not having the drawbacks exposed here above.

Furthermore such composition must allow the identification of the active components present in the preparation. The heat-killed preparations which
20 include heating at 120°C during 10 to 30 minutes would never allow identification of heat labile components.

Accordingly it is one object of the present invention to provide a novel killed Gram positive bacterial preparation such as killed Gram positive facultative intracellular bacterial preparation, for example a killed mycobacterial preparation
25 and compositions containing it, useful in the treatment of diseases comprising an immune dysregulation such as a Th1-Th2 imbalance, for example cancer, autoimmune diseases and allergic disorders without important adverse reactions.

It is another object of the present invention to provide a method of preparing said killed Gram positive bacterial preparation which besides its capacity for
30 the treatment of diseases including immune dysregulations, also preserves the struc-

ture of the molecules from the bacteria cells in order to allow the identification of the active molecules from these bacterial cells.

Surprisingly, the Inventors have shown that Gram positive bacteria such as Gram positive facultative intracellular bacteria, for example mycobacteria which are killed by "soft methods" which do not denature the molecules from the bacteria cells are able to stimulate leukocytic regulatory cells (CD4+ CD25+ T cells and/or B cells and/or dendritic cells) *in vivo* when they are administered to subjects suffering from asthma, or other immune dysregulation. The stimulation of these regulatory cells produces a reshaping of the immune reactivity of the asthmatic subjects which has a local and a systemic effect for a prolonged period (several weeks); as a result, the allergic subjects treated with these killed Gram positive intracellular bacterial preparations are protected from asthma for a prolonged period.

The Inventors have also shown that these killed bacterial preparations wherein the structure of the molecules from the bacteria cells is preserved, do not induce adverse effects such as inflammation, anemia, thrombopenia and induction of TNF- α production.

The present invention relates to a bacterial preparation, characterized in that:

- it contains killed Gram positive bacteria, obtainable by a process which does not denature the structure of the molecules from the bacteria cells, and
- it is able to induce, *in vivo*, a modulation of the immune response against an antigen (immunomodulatory preparation).

Immunomodulatory preparation means a preparation which is able to modify the ratio between immune regulatory and immune helper cells before or after the induction of an immune response against any antigen (self antigen or foreign antigen). Immune regulatory cells include leukocytic regulatory cells such as CD4+ CD25+ T cells and/or B cells and/or dendritic cells. For example, CD4+ CD25+ T cells are described in Schevach et al., *Nat. Rev. Immunol.*, 2002, 2, 389-400.

According to an advantageous embodiment of said bacterial preparation, it contains killed Gram positive facultative intracellular bacteria.

Gram positive facultative intracellular bacteria means Gram positive bacteria with the capacity of growing in synthetic medium *in vitro* as well as of infecting eukaryotic cells from a mammalian or non-mammalian host, *in vivo* and multiplying in those cells, for example macrophages.

5 According to an advantageous embodiment of said bacterial preparation, it contains killed Gram positive facultative intracellular bacteria chosen from *Listeria sp.*, *Corynebacterium sp.*, and Actinomycetes comprising *Mycobacteria sp.*, *Nocardia sp.* and *Rhodococcus sp.*.

 Preferably said bacterial preparation contains *Mycobacteria bovis*,
10 more preferably *Mycobacteria bovis* BCG.

 A process which does not denature the structure of the molecules from the bacteria cells means a process which results in no extensive denaturation of the spatial configuration of the molecules; preferably, said process preserves the three-dimensional structure of the macromolecules from the bacteria cells such as proteins,
15 polysaccharides and lipids.

 These processes which are denominated "soft processes" include with no limitation the use of physical means which disrupt the bacteria cell membranes while preserving the structure of its macromolecular components. These processes include with no limitation: extended freeze-drying, grinding in the presence
20 of silica or zirconium beads, use of the so-called "French press", sonication and gamma-rays irradiation. Other processes which may be used for obtaining the killed bacterial preparation as defined above are known to those of ordinary skill in the art.

 - For example, extended freeze-dried killed bacterial preparation means that essentially all the water has been removed from said preparation; thus, the
25 extended freeze-dried killed bacterial preparation contains less than 1.5% of residual water, preferably less than 1 % and more preferably less than 0.5%. However, in non-optimal freeze-drying conditions, when the preparations of freeze-dried bacteria contain more residual water (about 10 %), i.e. all the bacteria are not killed, killing of the residual living bacteria is alternatively obtained by contacting said preparations
30 with air (atmospheric pressure); such preparations have the same properties and activity as the above described extended freeze-dried killed bacterial preparations. The

residual water in the extended freeze-dried killed bacterial preparation is for instance determined by the coulometric method of Karl Fisher.

The absence of denaturation of the molecules from the killed Gram positive bacterial preparation according to the invention are verified by any method well-known in the art. For example, the structure of the proteins can be verified by gel electrophoresis of the killed Gram positive bacterial preparation extracts, in denaturing and non-denaturing conditions according to Laemmli, *Nature*, 1970, 277, 680-, by comparison with protein extracts obtained from living bacteria; the proteins are visualised either directly, by staining of the gel with an appropriate dye or after transfer of the proteins onto membranes and staining with appropriate antibodies directed to the Gram positive bacteria proteins. Other methods such as gel filtration or mass spectrometry can also be used to verify the structure of purified molecules from the killed Gram positive bacterial preparation according to the invention.

According to an other advantageous embodiment of said bacterial preparation, it contains extended-freeze-dried killed bacteria obtainable by an extended freeze-drying process.

Preferably, said extended-freeze-dried killed bacterial preparation is prepared by:

- (i) harvesting a culture of live bacteria cells,
- (ii) washing the bacteria cells in water or in an aqueous solution of a salt such as borate,
- (iii) freezing the bacteria cells in water or in an aqueous solution of a salt such as borate,
- iv) killing the frozen bacteria cells by drying them in a lyophiliser, for a time sufficient to remove at least 98.5 % of the water, preferably at least 99 % and more preferably at least 99.5%, and
- (v) collecting the extended freeze-dried killed bacteria cells.

Alternatively, said extended-freeze-dried killed bacterial preparation is prepared by a process as defined above with the exception that the washing of the bacteria cells in (ii) is omitted.

Preferably, step (iv) is performed at a drying chamber pressure of about 0.02 mBar to 0.2 mBar; more preferably 0.06 mBar to 0.1 mBar, during at least 10 to 12 hours, more preferably over several days with a low heat input and a low cold vapor trap temperature to ensure low temperature of dried killed bacterial preparation throughout the drying (no denaturation of the bacterial cells molecules).

In the freeze-drying process, the extended freeze-drying state is reached when in the absence of air leaks in the freeze-dryer, the vacuum-chamber pressure shows no more variation when the ice condenser and the vacuum pump are separated from the vacuum chamber for some seconds ; this means that no more water is capable of being removed from the freeze-dried bacteria (=stable water vapour pressure). For instance, when the cold trap is at -52°C, the pressure in the drying chamber may be in the range of 0.06-0.120 mBar, usually of about 0.09 mBar, for obtaining effectively extended freeze-dried killed bacteria.

The invention relates also to different fractions of the killed bacterial preparation according to the invention, selected in the group consisting of:

- a fraction A consisting of an organic solvent extract of said killed bacterial preparation, to eliminate phospholipids,
- a fraction B consisting of a glycosidase-treated extract of said killed bacterial preparation, to eliminate glyco-derived components, such as peptidoglycans,
- a fraction C consisting of a DNase and/or a RNase-digested extract of said killed bacterial preparation, to eliminate nucleic acids,
- a fraction D, consisting of a protease-treated extract of said killed bacterial preparation, to eliminate proteins,
- a fraction E, consisting of a extract of said killed bacterial preparation successively treated by an organic solvent, a glycosidase, a DNase and/or a RNase, and finally a protease.

Preferably:

- said fraction A is obtained by extraction with a methanol/chloroform mixture.
- said fraction B is obtained by digestion with lysozyme.

- said fraction C is obtained by digestion with DNase I and/or RNase

A.

- said fraction D is obtained by digestion with subtilisin.

According to an advantageous embodiment of said fractions they are
5 isolated from an extended freeze-dried bacterial preparation, as defined above.

According to an advantageous embodiment of said fractions, they consist of mycobacteria fractions, preferably *Mycobacteria bovis* fractions, more preferably *Mycobacteria bovis* BCG fractions.

The invention also relates to a pharmaceutical composition for the
10 prevention or the treatment of diseases comprising an immune dysregulation such as a Th1-Th2 imbalance, comprising an effective amount of killed bacterial preparation and/or at least one fraction thereof, as defined above, a pharmaceutically acceptable carrier and/or an additive, and/or an adjuvant, and/or an immunostimulant and/or an immunomodulator distinct from the bacterial preparation according to the invention.

15 Adjuvant means a natural or synthetic product which potentiates the specific immune response to an antigen (antibody production, B and T cell activation) when administered in association with said antigen.

Immunostimulant means a natural or synthetic product which induces a non-specific immune response when administered in association with an
20 antigen (for example an increase of the phagocytosis of the antigen).

According to the invention, said composition may advantageously be used for treating cancers, autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, Crohn disease and diabetes mellitus, and allergic disorders such as asthma, allergic rhinitis and atopic dermatitis.

25 According to an advantageous embodiment of said composition, it consists of mycobacteria, preferably *Mycobacteria bovis*, more preferably *Mycobacteria bovis* BCG.

According to another advantageous embodiment of said composition, it consists of extended-freeze-dried killed bacteria obtainable by an extended
30 freeze-drying process as defined above; preferably said extended-freeze-dried killed bacteria are obtained by a freeze-drying process as defined above.

According to yet another advantageous embodiment of said composition, it is in a form suitable to be administered by the intranasal route.

According to yet another advantageous embodiment of said composition, it is in a form suitable to be administered by the oral or sublingual routes.

5 In general, the composition may be administered by parenteral injection (e.g., intradermal, intramuscular, intravenous or subcutaneous), intranasally (e.g. by aspiration or nebulization), orally, sublingually, or topically, through the skin or through the rectum.

As specified, in one embodiment, the composition of the present
10 invention is in a form suitable for delivery to the bronchopulmonary mucosal surfaces. For example, the composition may be suspended in a liquid formulation for delivery to a patient in an aerosol form or by means of a nebuliser device similar to those currently employed in the treatment of asthma.

As specified, in another embodiment, the composition of the present
15 invention is in a form suitable for oral administration. For example, the composition may be in the form of tablets, ordinary capsules, gelatin capsules or syrup for oral administration. These gelatin capsule, ordinary capsule and tablet forms can contain excipients conventionally used in pharmaceutical formulation, such as adjuvants or binders like starches, gums and gelatin, adjuvants like calcium phosphate, disintegrating agents like cornstarch or alginic acids, a lubricant like magnesium stearate,
20 sweeteners or flavourings. Solutions or suspensions can be prepared in aqueous or non-aqueous media by the addition of pharmacologically compatible solvents. These include glycols, polyglycols, propylene glycols, polyglycol ether, DMSO and ethanol.

The composition may additionally contain a pharmaceutically
25 acceptable carrier and/or an additive and/or an immunostimulant and/or an adjuvant such as a liposome containing the bacteria cells or fraction(s) thereof according to the present invention; said one or more additives used for preparing pharmaceutical compositions may be chosen among antiaggregating agents, antioxidants, dyes, flavor enhancers, or smoothing, assembling or isolating agents, and in general among any
30 excipient conventionally used in the pharmaceutical industry.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical composition of the present invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline
5 buffer, lactose, glutamate, a fat or a wax. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g. polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable
10 biodegradable microspheres are disclosed, for example in US Patent 4,897,268 and 5,075,109.

Any of the variety of adjuvants may be employed in the compositions of the present invention to enhance the immune response. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism or to create
15 controlled inflammatory reactions, such as aluminium hydroxide or mineral oil, and a non-specific stimulator of immune response, such as lipid A, *Bordetella pertussis* toxin. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Freund's complete adjuvant which can not be used for injection in human. Other suitable adjuvants which can be used in human include
20 aluminium hydroxide, biodegradable microspheres, monophosphoryl A and Quil A.

The preferred frequency of administration and effective dosage will vary from one species and one subject to another. For example, the amount of extended freeze-dried killed mycobacterial preparation or fraction(s) thereof, is in a dose ranges which is equivalent in mice to about 1 μg to 10000 μg material (approximately 10^6 to 10^{10} CFU); preferably from 10 μg to about 1000 μg ; more preferably
25 from about 10 μg to 100 μg . The doses may be higher or lower, depending on the body surface of the species or the subjects and on the frequency of administration. For example, the administration of one or two doses once every two months may be more efficient in the treatment of asthma.

30 The invention also relates to products containing a bacterial preparation according to the invention or fractions thereof and a product selected from

the group consisting of anticancer, anti-diabetes and immunomodulatory drugs, as a combined preparation for simultaneous, separate or sequential use in the prevention and/or the treatment of diseases comprising an immune dysregulation.

Preferably, said products are chosen from anti-histaminic and anti-inflammatory drugs.

The invention also relates to the use of a bacterial preparation according to the invention or fractions thereof for the preparation of a medicament for the treatment of asthma, to be administered by the oral, sublingual, parenteral or intranasal route, in a dose range which is equivalent in mice to 1 μ g to 10 000 μ g, preferably 10 μ g to 1 000 μ g, most preferably 10 μ g to 100 μ g, at two months intervals, starting from at least two weeks before the usual period of allergen exposure.

The invention also relates to a use of a bacterial preparation according to the invention or fractions thereof for the preparation of a medicament for the prevention of asthma in human baby, to be administered by the oral, sublingual, parenteral or intranasal route, in a dose range which is equivalent in mice to 1 μ g to 10 000 μ g, preferably 10 μ g to 1 000 μ g, most preferably 10 μ g to 100 μ g.

The instant invention also relates to a method for the preparation of said extended freeze-dried killed bacterial preparation, characterized in that it comprises at least the steps of:

- (i) harvesting a culture of live bacteria cells,
- (ii) washing the bacteria cells in water or in an aqueous salt such as borate,
- (iii) freezing the bacteria cells in water or in an aqueous salt such as borate,
- (iv) killing the frozen bacteria cells by drying them in a lyophiliser, for a time sufficient to remove at least 98.5% of the water, preferably at least 99 % and more preferably at least 99.5%, and
- (v) collecting the extended freeze-dried killed bacteria cells.

An alternative method for the preparation of said extended freeze-dried killed bacterial preparation, comprises steps (i), (iii), (iv) and (v); the washing of the bacteria cells in step (ii) is omitted.

Preferably, step (iv) is performed at a drying chamber pressure of about 0.02 mBar to 0.2 mBar; more preferably 0.06 mBar to 0.1 mBar, during at least 10 to 12 hours, in order to remove at least 98.5% of the water while preventing melting of the frozen bacteria and maintaining the dried killed bacteria below the
5 bacterial cells molecules denaturation temperature.

According to a advantageous embodiment of said method, it consists of the preparation of extended freeze-dried killed mycobacteria.

The instant invention also relates to the use of said killed bacteria cells preparation or fraction(s) thereof for the preparation of a medicament for the
10 prevention and/or the treatment of diseases comprising an immune dysregulation such as a Th1-Th2 imbalance, e.g. cancers, autoimmune diseases such as multiple sclerosis, diabetes mellitus and Crohn disease, allergic disorders, such as asthma, allergic rhinitis or atopic dermatitis.

According to said use, said killed bacterial preparation or fraction(s)
15 thereof are associated with a pharmaceutically acceptable carrier, and/or an immunostimulant, and/or an adjuvant and/or any conventional additives as defined hereabove.

According to an advantageous embodiment of the invention mycobacteria, preferably *Mycobacteria bovis*, more preferably *Mycobacteria bovis* BCG are
20 used.

According to another advantageous embodiment of the invention, extended-freeze-dried killed bacteria obtainable by an extended freeze-drying process are used.

The killed bacterial preparations according to the invention have the
25 following advantages:

- they protect the subjects against diseases comprising an immune dysregulation such as a Th1-Th2 imbalance, e.g. asthma for example,
- they are not infectious and may be administered to immunodeficient subjects,
- 30 - they do not lead to inflammatory adverse effects, probably due in particular to the fact that they do not induce the production of TNF- α , which can lead

to toxicity and necrotizing activity. In addition, they do not produce other side effects including anemia and thrombopenia,

- they can be used without induction of a delayed type hypersensitivity to BCG purified protein derivatives (PPD) and thus do not interfere with the
5 diagnosis of tuberculosis,

- and they are prepared by a method that preserves the structure of the molecules from the bacteria cells (no extensive denaturation of the molecules), and thus allows the identification of the immunomodulatory molecules from said killed bacterial preparations.

10 The present invention also relates to a kit for the treatment of a disease comprising an immune dysregulation, characterised in that it comprises at least a pharmaceutical composition according to the invention, means of administration of said pharmaceutical composition and/or means for insuring compliance with the prescribed treatment.

15 The present invention will be further illustrated by the additional description and drawings which follows, which refers to examples illustrating the protective effect of extended freeze-dried killed BCG on asthma in mice, the analysis of the immune response induced by extended freeze-dried killed BCG, heat-killed BCG and living BCG, or of the production of extended freeze-dried killed BCG and
20 fractions thereof. It should be understood however that these examples are given only by way of illustration of the invention and do not constitute in anyway a limitation thereof.

- figure 1 illustrates the standard method of preparation of freeze-dried living BCG vaccine (A) by comparison with the method of preparation of
25 extended freeze-dried killed BCG according to the present invention (B): the pressure in mBar (-Δ-) and the temperature in degree Celsius of the samples (-O-) or of the trays (-●-) are indicated over the time in hours.

- figure 2 illustrates the protective effect of extended freeze-dried killed BCG (Lyoph killed BCG) on broncho-pulmonary hyper-reactivity in a particularly stringent (highly
30 Th2) BP2 mice model of asthma as assayed by the reduction in ventilation. The BP2 strain of mice is highly responsive along the Th2/asthma axis. The mice are also

sensitized with OVA in combination with alum, known to be a Th2 stimulant. By comparison, in this stringent mice model, no significant effect is observed in the groups treated with living BCG or heat-killed BCG. The area under the curve (cm^2) is presented: individual (-O-), mean \pm SEM per group of mice (-●-).

5 - figure 3 illustrates the protective effect of extended freeze-dried killed BCG (Lyoph killed BCG) on broncho-pulmonary hyper-reactivity in the BP2 mice model of asthma using OVA as allergen, as assayed by broncho-pulmonary resistance. By comparison, no significant effect is observed in the groups treated with living BCG or heat-killed BCG. Enhanced pause (Penh) measured at 1 minute interval
10 are presented: individual data (-O-), mean \pm SEM per group of mice (-●-).
***Statistically significant ($p < 0.001$) protective effect on broncho-pulmonary hyper-reactivity in the extended freeze-dried-killed BCG treated group.

 - figure 4 illustrates the protective effect of extended freeze-dried killed BCG (Lyoph killed BCG) against the increase of eosinophils in the lungs, in the
15 BP2 mice model of asthma, as assessed by cell numeration of the bronchoalveolar lavages (BAL). By comparison, no significant effect is observed in the groups treated with living BCG or heat-killed BCG. Individual data (-O-) and mean \pm SEM per group of mice (-●-) are presented.

 - figure 5 illustrates the protective effect of extended freeze-dried
20 killed BCG (Lyoph killed BCG) against the increase of neutrophils in the lungs, in the BP2 mice model of asthma, as assessed by cell numeration of the bronchoalveolar lavages. By comparison, no significant effect is observed in the mice treated with living BCG or heat-killed BCG. Individual data (-O-) and mean \pm SEM per group of mice (-●-) are presented.** Statistically significant reduction of the number of neutro-
25 phils ($p < 0.01$) in the extended freeze-dried killed BCG treated group.

 - figure 6 illustrates the advantageous absence of lung inflammation or tissue damage, with extended freeze-dried killed BCG (Lyoph killed BCG) in the BP2 mice model of asthma, as assayed by the very low level of fibronectin in bronchoalveolar lavages. By comparison, a significantly higher level of fibronectin in
30 the BAL is observed in the groups treated with living BCG or heat-killed BCG. Data

are expressed in IU/ml; individual data (-O-), mean \pm SEM per group of mice (-●-).*** Statistically significant reduction of the level of fibronectin ($p < 0.001$) in the extended freeze-dried killed BCG treated group.

5 - figure 7 illustrates the level of IL-5 in bronchoalveolar lavages from groups of mice that have been treated with living BCG, heat-killed BCG, extended freeze-dried killed BCG (Lyoph killed BCG), or non-treated. Data are expressed in pg/ml; individual data (-O-), mean \pm SEM per group of mice (-●-).

10 - figure 8 illustrates the level of IL-5 in the serum from groups of mice that have been treated with living BCG, heat-killed BCG, extended freeze-dried killed BCG (Lyoph killed BCG), or non-treated. Data are expressed in pg/ml; individual data (-O-), mean \pm SEM per group of mice (-●-).

15 - figure 9 illustrates the level of IFN- γ (pg/ml) in lung cultures from groups of mice that have been treated with living BCG, heat-killed BCG, extended freeze-dried killed BCG (Lyoph killed BCG), or non-treated; the cultures have been stimulated *in vitro* with purified BCG secreted protein (A) or with anti-CD3 antibodies (B). Individual data (-O-) and mean \pm SEM per group of mice (-●-) are presented.

20 - figure 10 illustrates the level of OVA-specific IgE in the serum of mice that have been treated in a curative protocol with living BCG, heat-killed BCG, extended freeze-dried killed BCG (Lyoph killed BCG) or non-treated. Individual data (-O-) and mean \pm SEM per group of mice (-●-) are presented.

25 - figures 11A and 12A illustrate the production of IL-12 (p70 and p40) by alveolar macrophages from respectively BP2 and BALB/c mice immunised with OVA, that have been stimulated *in vitro* by either heat-killed BCG (Heated) (-◆-), or extended freeze-dried killed BCG (Lyophilized) (....■....). Data are expressed in pg/ml per 100 000 cells.

- figures 11B (BP2) and 12B (BALB/c) illustrate the absence of production of TNF- α in the macrophages from mice stimulated by extended freeze-dried killed (Lyophilized) compared to significant level of TNF- α produced in the

macrophages from mice stimulated by heat-killed BCG (Heated). Data are expressed in or ng/ml per 100 000 cells.

- figure 13 illustrates the absence of delayed type hypersensitivity (DTH) to BCG purified proteins derivatives (PPD) in mice that have been treated with
5 extended freeze-dried killed BCG (Lyoph killed BCG). By comparison DTH is observed in mice that have been treated either with living BCG or with heat-killed BCG. Individual data (-O-) and mean \pm SEM per group of mice (-●-) are presented.
*** Statistically significant absence of foot-pad swelling ($p < 0.001$) in the extended freeze-dried killed BCG treated group.

10 - figures 14 illustrate the protective effect of extended freeze-dried killed BCG (EFD-BCG) on broncho-pulmonary hyper-reactivity to histamine in a guinea-pig model of asthma. Groups of OVA immunised animals were treated with 10 μ g (figure 14B) or 100 μ g (figure 14C) of extended freeze-dried killed BCG, or non-treated (figure 14A). Concentration of histamine able to create broncho-constriction
15 before (black symbols) and after (white symbols) aerosol administration of OVA are presented for each animal. *** Statistically significant protective effect in the group treated with 10 μ g ($p < 0.01$) or 100 μ g ($p < 0.001$) extended freeze-dried killed BCG (Kruskal –Wallis non parametric test).

- figure 15A illustrates the protective effect of extended freeze-dried
20 killed BCG (EFD1) against the increase of eosinophils in the lungs, in the BP2 mice model of asthma, as assessed by cell numeration of the bronchoalveolar lavages. By comparison, no significant effect is observed in the groups treated with living BCG or heat-killed BCG. The data are expressed as mean \pm SEM per group of mice; $n = 7$ for each group. *Statistically significant reduction of the number of eosinophils ($p < 0.05$)
25 in the extended freeze-dried killed BCG treated group.

- figure 15B illustrates the level of IL-5 in bronchoalveolar lavages from groups of mice that have been treated with living BCG, heat-killed BCG, extended freeze-dried killed BCG (EFD1), or non-treated. The data are expressed as mean \pm SEM per group of mice; $n = 7$ for each group.

30 - figure 16 illustrates the preservation of the structure of the proteins in the extended freeze-dried killed BCG preparations as assessed by SDS-PAGE and

(A) staining with anti-mycobacteria antibodies after transfer onto PVDF membranes or
(B) staining with Aurodyne. The proteins appear as discrete bands without smear.

- figure 17 illustrates the protective effect of extended freeze-dried killed BCG (EFD1) on broncho-pulmonary hyper-reactivity in the BP2 mice model of asthma using a water soluble ray-grass pollen extract as allergen, as assayed by broncho-pulmonary hyper-reactivity. Enhanced pause (Penh) measured between 3 and 8 minutes after methacholine administration are presented: data are presented as mean \pm SEM per group of mice; n = 8 for each group. **Statistically significant ($p < 0.01$) protective effect on broncho-pulmonary hyper-reactivity in the extended freeze-dried-killed BCG treated group.

- figure 18A and B illustrates the protective effect of extended freeze-dried killed BCG (EFD1) against the increase of polymorphonuclears and eosinophils in the lungs, in the BP2 mice model of asthma, as assayed by cell numeration of the bronchoalveolar lavages. The data are expressed as mean \pm SEM per group of mice. Statistically significant reduction of polymorphonuclears number (** $p < 0.01$) in the groups treated with 1 mg and 10 mg extended freeze-dried killed BCG; statistically significant reduction of the eosinophils number: ** $p < 0.01$, * $p < 0.05$ and * $p < 0.05$ in the groups treated with respectively 100 μ g, 1 mg and 10 mg of extended freeze-dried killed BCG.

- figure 18C illustrates the correlation between the decrease of the number of polymorphonuclears and eosinophils in the lungs of the extended freeze-dried BCG (EFD1) treated groups (10 μ g, 100 μ g, 1 mg and 10 mg), and the decrease of IL-4 production as assessed by cells and cytokines analysis in the bronchoalveolar lavages. The data are expressed as mean \pm SEM per group of mice. Statistically significant reduction of the level of IL-4 (** $p < 0.01$) in the groups treated with 10 μ g, 100 μ g, 1 mg and 10 mg of extended freeze-dried killed BCG.

- figure 19 illustrates the prevention of leukocyte infiltration of the lungs in the extended freeze-dried killed BCG (EFD1) treated groups, in the BP2 mice model of asthma, as assayed by lungs cellular infiltrate numeration. The data are expressed as mean \pm SEM per group of mice; n = 5 for each group. Statistically

significant reduction of the number of cells (** $p < 0.001$) in the groups treated with 10 μ g extended freeze-dried killed BCG.

- figure 20 illustrates the prevention of leukocyte infiltration of the lungs in the extended freeze-dried killed BCG (EFD) treated groups, in the BP2 mice model of asthma, as assayed by flow cytometry analysis of the macrophages (CD11b+), the polymorphonuclear cells (Gr1+) and the dendritic cells (CD11c+). The data are expressed as mean \pm SEM per group of mice ; $n = 5$ for each group. Statistically significant reduction of the number of: (i) macrophages and dendritic cells in all the groups treated with extended freeze-dried killed BCG (10 μ g 100 μ g, 1 mg and 10 mg), and (ii) polymorphonuclears in the groups treated with 100 μ g, 1mg and 10 mg extended freeze-dried killed BCG only.

- figure 21 illustrates the level of IFN- γ and IL-10 (pg/lung) in lung cultures from groups of mice which have been treated with living BCG, heat-killed BCG, extended freeze-dried killed BCG (EFD1) or non-treated and challenged with OVA or not; the cultures have been stimulated *in vitro* with purified BCG secreted protein Data are expressed as mean \pm SEM per group of mice ; $n = 4$ for each group. Statistically significant increase of IL-10 production (***) $p < 0.001$ in the groups treated with extended freeze-dried killed BCG only.

- figure 22 illustrates the absence of anemia (A) and thrombopenia (B) after intravenous administration of extended freeze-dried killed BCG (EFD-BCG), contrary to administration of living BCG or heat-killed BCG by the same route.

- figure 23 illustrates the minimal inflammatory reaction at the site of injection after subcutaneous injection of extended freeze-dried killed BCG (EFD1), by comparison with living BCG and heat-killed BCG, as assessed by footpad increase measurement every week for 10 weeks after the injection. Arrow indicate the day of injection; D0 and D36 for EFD and heat-killed BCG, D0 only for living BCG.

- figure 24 illustrates the reduced TNF- α production after LPS stimulation of lung explants 18 days after intravenous injection of extended freeze-dried killed BCG (EFD1), by comparison with injection of living BCG and heat-killed BCG by the same route. No significant production of TNF- α (***) $p < 0.001$ in the extended freeze-dried killed BCG group compared to the untreated group. By

comparison significant production of TNF- α is observed in the living BCG and heat-killed BCG treated groups.

- figure 25 illustrates the increase of the CD11c⁺ Gr1⁺ B220⁺ plasmacytoid dendritic cells number in the spleen from BALB/c mice immunized 90 days previously with ovalbumine, treated or not with extended freeze-dried killed BCG (EFD1) at days 45 and 65 and challenged or not with ovalbumine. *** Statistically significant increase of the CD11c⁺ Gr1⁺ B220⁺ cells number ($p < 0.001$) in the treated group, no increase of the CD11c⁺ CD8 α ⁺ cells number in the treated group.

- figure 26 illustrates the increase of the CD4⁺ CD25⁺ IL-10⁺ cells number in the spleen from BALB/c mice immunized 90 days previously with ovalbumine, treated or not with extended freeze-dried killed BCG (EFD1) at days 45 and 65 and challenged or not with ovalbumine; spleen cells were stimulated *in vitro* with OVA or BCG culture supernatant or non-stimulated before analysis of the leukocyte populations. *** Statistically significant increase of the CD4⁺ CD25⁺ IL-10⁺ cells number ($p < 0.001$) in the treated groups.

- figure 27 illustrates the protective effect of extended freeze-dried killed BCG (EFD1) given by the oral route, in the mouse model of asthma, as assessed by lungs cell infiltrate numeration. *** Statistically significant decrease of the lungs cell number in the treated group ($p < 0.001$).

- figure 28 illustrates the protective effect of one dose of extended freeze-dried killed BCG (EFD) injected by the subcutaneous route, as assessed by the prevention of bronchopulmonary hyper-reactivity in the BALB/c model of asthma.

- figure 29 illustrates the delay of action of extended freeze-dried killed BCG (EFD) injected by the subcutaneous route, as assessed by the prevention of bronchopulmonary hyper-reactivity in the BP2 model of asthma. The prevention of bronchopulmonary hyper-reactivity occurs only if a delay of two to three weeks is present between extended freeze-dried killed BCG administration and challenge.

- figure 30 illustrates the duration of action of extended freeze-dried killed BCG (EFD1), injected by the subcutaneous route, as assessed by the prevention of bronchopulmonary hyper-reactivity in the BP2 model of asthma; the extended freeze-dried killed BCG protective effect persists at least for two months.

EXAMPLE 1: PREPARATION OF EXTENDED FREEZE-DRIED KILLED MYCOBACTERIA

1) Material and methods

The *Mycobacterium bovis* BCG cells (BCG Pasteur vaccine strain 1173P2, deposited at the Collection Nationale de Cultures de Microorganismes, 25 rue du Docteur Roux, 75724 PARIS Cedex 15 (France), on April 24, 1978, under n° I-059 (M. Gheorghiu and coll., 1983, Bull. Inst. Pasteur, 81, 281-288) are grown in sterile Sauton medium (HOOCCH(NH₂O)CH₂CONH₂H₂O (asparagine), 4 g/l; C₆H₈O₇·H₂O (citric acid), 2 g/l; K₂HPO₄ (di-potassium hydrogenophosphate), 0.5 g/l; MgSO₄·H₂O (magnesium sulphate), 0.50 g/l; FeIII citrate, 0.05 g/l; glycerol, 60 ml; Zinc sulphate solution (0.155 g Zinc sulphate in 10 ml pyrogen free water), 240 µl/l, pH 7). More precisely the 1173 P2 strain is grown in 250 ml spherical culture flask containing 130 ml sterile Sauton medium, at 37°C, for 14 days, corresponding to the time the culture ends its exponential phase.

The culture is then centrifuged at 2000 g for 10 minutes, at 4°C to pellet the BCG cells. The cell culture supernatant is discarded and the pellet is washed extensively (three times) in distilled water; each wash consists in resuspending said pellet in distilled water (20 volumes of water per volume of cell-pellet) and centrifuging the cells at 2000 g for 10 minutes, at 4°C.

After the last wash, the pellet is resuspended in a volume of distilled water equivalent or twice that of the volume of the pellet. 20 g of bacteria in 50 ml of water is layered and frozen on the wall of bottles, in order to form a layer of approximately 1 cm in thickness. The mixture is freeze-dried at -60°C.

The frozen suspension of cells in water is then extended freeze-dried under the conditions of the extended freeze-drying protocol described in figure 1B, to remove essentially all the water (residual water <1.5%). Briefly, the pressure is rapidly decreased to 0.150 mBar, the temperature of the ice condenser being approximately -50°C to -54°C, and the freeze-drying process begins. The samples remain frozen, the temperature being not exactly monitored. The pressure is then maintained at 0.1 mBar for 34 h, in order to remove essentially all the water (residual water <1.5%). There is a progressive increase up to room temperature (approximately 20°C). At the end of the

drying, the pressure is then slowly returned to atmospheric pressure with air, at room temperature (approximately 20°C) and the extended freeze-dried killed BCG cells (approximately 2 g) are harvested and stored at room temperature, under air atmospheric pressure.

5 The residual water present in the extended freeze-dried killed BCG preparation is determined using the coulometric method from Karl Fisher and a 756 KF Coulometer (METHROM), according to the manufacturer's instructions.

 The presence of viable cells in the extended freeze-dried killed preparation is verified by:

10 - determination of the number of colony forming units (CFU) per ml; 0.2 ml of a suspension of the extended freeze-dried killed preparation (1 mg, e.g. approximately 10^9 extended freeze-dried killed BCG cells in water) are plated on Middlebrook 7H10 agar medium (DIFCO), and the plates are incubated at 37°C for one month, and/or

15 - staining with the fluorescent dye CFDA-SE (Carboxyfluorescein diacetate- Succinimidyl Ester; MOLECULAR PROBES reference number C-1157) which stains viable cells only; 1 ml of a suspension of the extended freeze-dried killed preparation (approximately 10^8 extended freeze-dried killed BCG cells in water) are mixed with 100 µl of a 1/100 dilution in PBS of CFDA reagent (stock solution: 20 1mg/ml in DMSO). The mixture is incubated in the dark at room temperature for 60 min. Labeled bacteria are centrifuged at 3000 rpm for 15 minutes, washed twice in PBS and resuspended in the same buffer. Then, the presence of viable cells was examined by fluorescence microscopy.

2) Results

25 BCG cells were extended freeze-dried killed following the extended freeze-drying protocol described in material and methods and summarized in figure 1B. By comparison, BCG cells were freeze-dried following the standard method for preparation of live BCG vaccine: BCG cells (strain 1173P2) were cultured in Sauton medium at 37°C, harvested at the end of the exponential phase by centrifugation and 30 the BCG cell pellet was resuspended in sodium glutamate (1.5 g/100 ml H₂O), at the concentration of 4 mg/ml of BCG). The BCG suspension was distributed in vials (0.25

ml/vial), freeze-dried according to the protocol illustrated in figure 1A and each vial containing 1.5 ± 0.5 % H₂O was stored at room temperature under vacuum (0.06 mBar).

The viability of the BCG cells in the different preparations was
5 assayed as described in Material and Methods.

The following results were observed by the coulometric method of Karl Fisher, the culture assay and by the CFDA-SE staining test:

- 200 mg of extended freeze-dried killed BCG preparation contained
1.073 mg of water in a representative determination, which is equivalent to less than
10 0.5 % residual water.

- no living bacteria were detected in the extended freeze-dried killed BCG samples prepared according to the method of the invention,

- 50% to 60% living bacteria were detected in the freeze-dried killed BCG preparation according to the standard BCG vaccine preparation method.

15 **EXAMPLE 2: PREPARATION OF EXTENDED FREEZE-DRIED KILLED MYCOBACTERIA FRACTIONS**

1) Delipidated fraction (fraction A)

10 mg/ml of approximately 10^{10} extended freeze-dried killed BCG cells, suspended in borate buffer (Na₂B₄O₇ 10H₂O 0.363%; H₃BO₃ 0.525%; NaCl 0.619% and Tween 20 0.0005% in distilled water, pH 8), were centrifuged at 12 000 g
20 for 10 minutes and the supernatant was discarded. The pellet was resuspended in 1 ml of chloroform/methanol (9/1) for 24 h at room temperature to extract lipids. The chloroform/methanol is removed by centrifugation at 12 000 g per 10 min. The delipidated pellet from the chloroform/methanol extraction was vacuum-dried.

25 **2) Deglycosylated fraction (fraction B)**

10 mg/ml of approximately 10^{10} extended freeze-dried killed BCG cells, suspended in borate buffer (Na₂B₄O₇ 10H₂O 0.363%; H₃BO₃ 0.525%; NaCl 0.619% and Tween 20 0.0005% in distilled water, pH 8), were centrifuged at 12 000 g for 10 minutes and the supernatant was discarded. The pellet was resuspended in 1 ml
30 of 0.1 M Sodium acetate, for 24 h at 37°C, containing 1 mg lysozyme to remove the

peptidoglycan. The digestion product was heated 2 min at 96°C and the pellet was harvested by centrifugation for 1 h at 12 000 g.

3) DNase and RNase treated fraction (fraction C)

10 mg/ml of approximately 10^{10} extended freeze-dried killed BCG cells, suspended in borate buffer, were centrifuged at 12 000 g for 10 minutes and the supernatant was discarded. The pellet was resuspended in 1 ml of 40 mM Tris-HCl buffer containing 1 mg DNase and 1 mg RNase and 5 mM $MgCl_2$, for 24 h at 37°C, to remove the nucleic acids. The digestion product was heated 2 min at 96°C and the pellet was harvested by centrifugation for 1 h at 12 000 g.

4) Protease treated fraction (fraction D)

10 mg/ml of approximately 10^{10} extended freeze-dried killed BCG cells, suspended in borate buffer, were centrifuged at 12 000 g for 10 minutes and the supernatant was discarded. The pellet was resuspended in 1 ml of 0.1 M ammonium acetate buffer containing 0.1 mg subtilisin, for 23 h at 37°C. To remove the protein completely, 0.1 mg subtilisin was then added to the reaction mixture and incubated for 7 h at 37°C. The digestion product was heated 2 min at 96°C and the pellet was harvested by centrifugation for 1 h at 12 000 g.

5) Successive treatments (fraction E)

10 mg/ml of approximately 10^{10} extended freeze-dried killed BCG cells were centrifuged at 12 000 g for 10 minutes and the supernatant was discarded. The pellet was resuspended in borate buffer and treated successively with chloroform/methanol (9/1), lysozyme, DNase-RNase mixture and subtilisin as described above. The digestion product was heated 2 min at 96°C and the pellet was harvested by centrifugation for 1 h at 12 000 g.

6) Characterization of the protein and polysaccharide content of the extended freeze-dried killed BCG preparation.

a) Material and Methods

A preparation of extended freeze-dried killed BCG preparation (10 mg) was suspended in 1 ml of 4% butanol in water and distributed in two eppendorf tubes (500 μ l per tube) containing 1 g of zirconia/silica beads (0.1mm) (BIOSPEC PRODUCTS, INC.). The tubes were then placed in a Mixer Mills (MM301, RESCH)

at a frequency of 25 during 5, 15, 30, 60, 120, 180, 240, 300 or 470 minutes. The tubes were then centrifuged at 10000 rpm for 30 min; the supernatant was recovered, filtered (0.22µm filter, MILLIPORE) and the protein, the amino acids and the polysaccharide concentrations were determined as follows:

5 - total protein concentration was determined using Micro BCA Protein Reagent Kit (PIERCE),

 - amino-acids concentration determination and analysis was determined by mass spectrometry

 - total polysaccharide concentration was determined using anthrone
10 method, according to S. Melvin, *Anal. Biochem.* 1953, 25, 1656-.

 Aliquots (5 µg of proteins) of the bacteria extracts collected at 5, 60 or 300 min were separated by SDS-PAGE, the protein were then transferred onto a PVDF membrane and either stained with Aurodye® (Pharmacia) or with polyclonal antibodies directed to mycobacterial proteins.

15 b) Results

 The bacteria extracts collected after 5, 60 or 300 min of extraction contains respectively 280, 500 and 1258 µg of proteins ; 190, 420 and 880 µg of amino acids and 697, 1267 and 1765 µg of polysaccharides.

 The results presented in figure 16 show that the extended freeze-
20 drying process preserves the structure of the proteins which are present in the killed BCG preparations as indicated by the presence of discrete protein bands without smear ; the protein profile observed with the extended freeze-dried killed BCG preparation extracts is comparable to that obtained with living BCG preparations.

25 **EXAMPLE 3: PROTECTIVE EFFECT OF EXTENDED FREEZE-DRIED KILLED MYCOBACTERIA ON ASTHMA IN A MOUSE MODEL.**

1) Material and Methods

a) Animals

Male adult (6 to 7 weeks old) BP2 (H-2^d) mice were obtained from the Centre d'élevage R. Janvier (Le Genest, Saint Isle, France) and were maintained in animal
30 facilities in specific pathogen free-conditions.

b) Mouse model of the asthma-like allergen specific disease

Male adult BP2 mice were immunized by the subcutaneous route (in the dorsal part of the neck) with 1 µg ovalbumin (OVA; ICN Laboratories) and 1.6 mg aluminium hydroxide adjuvant in saline (0.4 ml final volume), on days 0 (D₀) and 7 (D₇). The
5 allergic response is provoked by intranasal challenge with 10 µg OVA in 50 µl saline on day 96-98. Alternatively, a water soluble ray-grass pollen extract is used as an allergen, in the same conditions as for OVA except that 10 µg is used for immunization.

c) Preparation of the mycobacterial immunomodulatory composition

10 Extended freeze-dried killed BCG, prepared as described in example 1 were tested as an asthma vaccine in the BP2 mouse, by comparison with the following BCG preparations :

- *Living BCG*

The *Mycobacterium bovis* BCG Pasteur vaccine strain 1173P2 was
15 grown as dispersed bacilli in Beck-Proskauer medium (Gheorghiu and coll., 1988, *J. Biol. Standard.*, 16, 15-26)) supplemented with 0.05% Triton WR 1339 (SIGMA) and 6% glucose. The bacteria were harvested at the exponential phase (5 to 7 days) and stored at -70°C in Beck-Proskauer medium supplemented with 0.05% Triton and 6% glycerol. The number of colony forming unit (CFU) per ml was determined by plating
20 suitable dilutions in phosphate-buffered saline (PBS) on Middlebrook 7H10 agar medium (DIFCO). The suspension was diluted at the concentration of 10⁸, 10⁹ or 10¹⁰ CFU/ml in PBS just before its injection (100 µl).

Heat-killed BCG

Living BCG prepared as above described was centrifuged at 3000
25 rpm for 15 minutes and the supernatant was discarded. The pellet was resuspended in saline, for example in a buffer containing borate (Na₂B₄O₇ 10H₂O 0.363%; H₃BO₃ 0.525%; NaCl 0.619% and Tween 20 0.0005% in distilled water, pH 8), and the suspension was autoclaved for 15 min at 115°C.

d) Curative protocol of administration of extended freeze-dried killed mycobacteria in the mouse model of asthma

Male adult BP2 mice are immunised with OVA as described above. Groups of 25 mice were injected with 100 µl of the BCG compositions prepared as described above.

- living BCG vaccine strain 1173P2 (10^7 CFU)
- heat-killed BCG (10^8 bacteria corpses which are equivalent to 10^7 CFU),
- extended freeze-dried killed BCG (10 µg to 10 mg; 10 µg is equivalent of 10^7 CFU), and
- PBS,

by the subcutaneous route (base of the tail), on D₄₂₋₄₅ and D₆₅₋₇₀ after the first immunisation with OVA.

The allergic response is provoked by intranasal challenge with 100 µg OVA in 50 µl saline on day 96-98 ; PBS alone is administered intranasally for comparison.

e) Broncho-pulmonary (BHR) analysis

A barometric plethysmographic equipment (BUXCO) that allows BHR study on unanesthetised animals was used. Animals immunized with OVA, treated with the different BCG preparations or non-treated were tested 24 h after the challenge with OVA or water soluble ray-grass pollen extract. The animals were placed in a plethysmographic chamber. Their basal ventilation parameters were recorded and during 20s or 1 minute they received inhalation of methacholin (ALDRICH), 100 mM in H₂O by using standard nebuliser, and their ventilation parameters were recorded during 10 minutes after methacholin aerosol administration. Reduction of the ventilation during the period is reported (area under the curve) and the bronchopulmonary resistance was expressed as enhanced pause (Penh), calculated as: (expiratory time(tE)/40% of relaxation time (trel) -1) x peak expiratory flow (PEF)/peak inspiratory flow (PIF) x 0.67, according to the manufacture' instructions. Every minute an average value of Penh was recorded. For the graphic representation, each value was expressed for every minute.

f) Bronchoalveolar lavage (BAL)

Mice were anesthetised by the intraperitoneal route with a lethal dose of urethane (1.5 g/kg; SIGMA). Circulating blood was removed from the abdominal aorte and a canula connected to a syringe was placed in the trachea, and the
5 lungs were washed three times with 0.7 ml PBS and the lavage was collected in a tube and kept on ice.

The number of cells present in the BAL was determined in an automatic counting equipment (ZBI), provided by COULTER. To determine the percentages of the different cell types (eosinophils, neutrophils, macrophages), the
10 BAL cells which have been cytospinned on glass slides, were stained with DIFF QUICK (BAXTER).

g) Fibronectin assay

BAL were centrifuged at 1000 rpm for 10 min at 4° C and the fibronectin present in the supernatant was assayed by competition Enzyme Immunometric
15 Assay (EIA), following standard protocols (Rennard and coll., Anal. Biochem., 1980, 205-214).

h) Preparation and analysis of pulmonary explants

The lungs were cut in small pieces of approximately 1 mm per 3 mm. Four or five pieces from each lung were placed in a well of a tissue culture
20 plate containing 1 ml of AIM V culture medium (Life Technologies). The pulmonary explants were cultured in medium alone for basal level determination; alternatively BCG culture filtrate (10 µg/ml) was added to the culture to reveal mycobacterial specific cells, or anti-CD3 monoclonal antibody was added to the culture to reveal reactivity of T lymphocytes. 24 h later, the presence of different lymphokines (IFN-γ,
25 TNF-α, IL-10...) was assayed by ELISA using commercial kits, according to the manufacturer's instructions.

The entire lung (without trachea) was dissociated in collagenase and DNase medium to enumerate the infiltrating cells on a Flow cytometer using various monoclonal antibodies labelled with different fluorochromes.

i) Histopathological analysis of the lungs

The entire lungs were fixed in buffered formaldehyde and processed for histopathological analysis. The Schiff staining was used to stain the mucus and the mucus containing cells. The hematoxy-eosin stain was used as a counter staining. The slides were examined and an evaluation of cell infiltration and mucus content was performed by one observer who scored the slide without knowledge of the origin of the sample (blind assay).

j) Cytokines assay

The cytokines which are either present in the serum and the BAL samples from the groups of immunised mice, or secreted by the lung explants cultures from these mice, were assayed as follows: IFN- γ , IL-10 and IL-12 were tested by ELISA; TNF- α and IL-5 were tested by EIA (Enzyme Immunometric Assay).

k) OVA-specific IgE assay

The level of OVA-specific IgE present in the serum was assayed by ELISA, following standard protocols (Hansen and coll., J. Immunol., 2000, 223-230).

l) Statistical analysis

The number (n) of animals per group was $4 < n < 8$. The student's two-tailed (t) test was performed for independent events.

2) Results

a) Broncho-pulmonary (BHR) analysis

Animals immunised with OVA or a water soluble ray-grass pollen extract and treated with the different BCG preparations, or non-treated, were tested 24 h after the challenge with OVA or ray-grass pollen.

The model chosen in these studies, using BP2 mice with Th2 background which are immunised with ovalbumine associated with aluminium hydroxide, a more potent adjuvant of allergic reaction than the freund incomplet adjuvant, is thus more stringent than the model of BALB/c mice immunized with ovalbumine in freund incomplete adjuvant used by CC Wand and coll., precited.

The data presented in figures 2 and 3 indicate that in this stringent model of asthma, extended freeze-dried killed BCG (10 μ g) has a statistically significant ($p < 0.001$) protective effect on broncho-pulmonary , as expressed by a lower

reactivity to metacholine compared to non-treated controls. By contrast, in this stringent model of asthma, no significant protective effect was observed in the animals treated with live BCG (10^7 CFU) or heat-killed BCG (10^8 bacterial corpses). A similar statistically protective effect ($p < 0.01$) on asthma is observed with the extended
5 freeze-dried killed BCG (EFD) preparation, when a water soluble ray-grass pollen extract is used as an allergen (figure 17).

b) Bronchoalveolar lavage (BAL)

A decrease of the number of eosinophils ($p < 0.05$) and neutrophils ($p < 0.05$) is observed in the groups of mice treated with 10 μ g, 100 μ g, 1 mg or 10 mg
10 of extended freeze-dried killed BCG (figures 4, 5, 15A, 18A and 18B).

The level of fibronectin present in airways exudation liquid which reflects the inflammatory response, shows a very significant decrease ($p < 0.001$) in the group treated with extended freeze-dried killed BCG (10 μ g, figure 6).

Assays as regards IL-5, a lymphokine implicated in the increase of
15 the number of eosinophils during allergic responses, show a decrease both in BAL and in the blood, in the group treated with extended freeze-dried killed BCG (figures 7, 8 and 15).

Assays as regards IL-4, a Th2 cytokine, shows a statistically significant decrease in BAL from the groups of mice treated with 10 μ g, 100 μ g, 1 mg or 10
20 mg of extended freeze-dried killed BCG (figure 18C).

By comparison, no significant effect was observed in mice treated with either living BCG (10^7 CFU) or heat-killed BCG (10^8 bacterial corpses).

These results show that extended freeze-dried killed BCG can induce an immune response capable of treating the symptoms of asthma as demonstrated by the decrease in the inflammatory response of the lung (production of eosinophils and neutrophils, exudation of fibronectin). In this stringent model of asthma
25 (BP2 mice) with Th2 background, no protection was observed in either living BCG or heat-killed BCG treated groups.

c) Analysis of the histopathology and the infiltrating cells of the lungs

The histopathological analysis of the lungs shows that the treatment with extended freeze-dried killed BCG (EFD) prevents leukocyte infiltration of the lungs and mucosal metaplasia of bronchial epithelia.

These results are confirmed by the flow cytometry analysis of the cells present in the lung tissues which demonstrates that the treatment with extended freeze-dried killed BCG induces a statistically significant ($p < 0.001$) decrease of leukocyte infiltration in the lungs (figure 19) ; analysis of the different leukocyte populations (figure 20), shows a decrease of the number of macrophages ($\text{CD}11\text{b}^+$), polymorphonuclear cells ($\text{Gr}1^+$) and dendritic cells ($\text{CD}11\text{c}^+$) in the lungs of the extended freeze-dried killed BCG treated groups.

d) Cytokine analysis

Lung biopsies from the groups of mice that have been treated with the different BCG preparations were cultured in the presence or in the absence of non-purified proteins secreted by BCG or anti-CD3 mAb, and the production of IFN- γ and IL-10 was assayed in the supernatant

The results presented in figures 9 and 21 indicate that IFN- γ production requires an immunisation with BCG (figure 9 A). The lower level IFN- γ observed in the extended freeze-dried killed BCG group (figure 9A) is not statistically significant from the level observed in the other BCG immunised groups. Non-specific stimulation of T lymphocytes with anti-CD3 antibodies shows similar levels of IFN- γ in the different groups (treated or non-treated with BCG) suggesting that immunisation with BCG does not modify the number of T lymphocytes present in the lungs (Figure 9B).

25 The results presented in figure 21 show also a statistically significant
($p < 0.001$) higher level of IL-10 in the extended freeze-dried killed BCG treated
groups compared to living BCG and heat-inactivated BCG treated groups.

The differences of the cytokines expression profile in the lungs between the extended freeze-dried killed BCG treated groups and the groups treated with the other BCG preparations indicate that extended freeze-dried killed BCG activity is correlated with modifications in the leukocyte populations present in the

lung tissues after allergen delivery. The high amount of IL-10 and low amount of IFN- γ released by lung explants could assert the presence of regulatory cells in lung tissues of the extended freeze-dried killed BCG groups, after allergen delivery.

e) Level of anti-OVA specific IgE

5 The immunization with OVA having been realized before treating with the different BCG preparations (curative protocol), the level of anti-OVA IgE is identical for all the groups (figure 10), suggesting that the protective effect observed with extended freeze-dried killed BCG does not involve a direct effect on IgE concentration.

10 **EXAMPLE 4: ANALYSIS OF THE CYTOKINES PRODUCED BY ALVEOLAR MACROPHAGES STIMULATED BY HEAT-KILLED OR EXTENDED FREEZE-DRIED KILLED BCG**

1) Material and methods

 BP2 and BALB/c mice (H-2^d) mice, obtained and maintained as
15 mentioned in example 3, were immunized with OVA as described in example 3b. One week after the last injection with OVA, mice were anesthetised by the intraperitoneal route with a lethal dose of urethane (1.5 g/kg; SIGMA). Circulating blood was removed from the abdominal aorte and a canula connected to a syringe was placed in the trachea, and the lungs were washed twice with 0.5 ml and 5 times with 1 ml of
20 HBSS medium (Life Technologies) containing 2% of foetal calf serum. The number of macrophages present in the BAL was determined in an automatic counting equipment (ZBI), provided by COULTER. The macrophages were centrifuged at 1000 rpm for 20 min at room temperature and their concentration was adjusted to 340. 10³ cells/ml in RPMI 1640 medium containing 3 % foetal calf serum, 2 mM glutamine, 100 UI/ml
25 penicilline and 10 µg/ml streptomycine. The macrophages were then distributed in a 96 well plate for tissue-culture (100 000 cells /well) and incubated for at least 2 hours at 37°C in the presence of 5% CO₂. After, removal of the non-adherent cells by several washes with RPMI medium, macrophages were cultured in the presence of the heat-killed BCG or extended freeze-dried killed BCG (5 equivalent CFU per macrophage
30 e.g. 1.6 10⁶ CFU/ml for both BCG preparations); *E. coli* lipopolysaccharide (LPS; 1 µl/ml) and OVA (10 µg/ml) were used as controls. The cell culture supernatant were

harvested 2, 4, 6, 8, 12, 24, 48, 72 and 96 hours after addition of the different preparations and frozen immediately at -20°C.

2) Results

The results presented in figures 11 and 12 show that:

5 1) IL-12 which is involved in the maturation of the Th1 cells and the production of anti-allergic cytokines (IFN- γ) is produced by BALB/c and BP2 mice alveolar macrophages stimulated by extended freeze-dried killed and heat-killed BCG preparations,

2) TNF- α a cytokine that can be associated with important inflammatory side-effects (necrosis, ulceration...) is produced only by BALB/c and BP2 mice alveolar macrophages stimulated by heat-killed BCG preparations; no TNF- α is produced by alveolar macrophages stimulated by extended freeze-dried killed BCG preparations at the concentrations used in the present experiments.

15 These results support the minimal adverse effects observed after extended freeze-dried BCG injection, contrary to the side effects observed after heat-killed and living BCG injection.

EXAMPLE 5: THE ADMINISTRATION OF EXTENDED FREEZE-DRIED KILLED BCG DOES NOT INDUCE DELAYED TYPE HYPERSENSITIVITY (DTH) TO BCG PURIFIED PROTEINS DERIVATIVES (PPD) .

20 1) Materials and methods

Mice were immunised with OVA and treated with different preparations of BCG, or non-treated, as described in example 3d. 100 days after the first subcutaneous injection with the different BCG preparations, the delayed type hypersensitivity (DTH) to BCG purified proteins derivatives (PPD) which is used for the diagnosis of tuberculosis in humans was performed on the different groups of mice. More precisely, 50 μ l of PPD in saline (4 μ g) was injected into the mice footpad and 24h later the footpad swelling was measured.

2) Results

30 The results presented in figure 13 show that mice immunised with extended freeze-dried killed BCG (equivalent to 10^7 CFU or 10 μ g) do not develop a delayed type hypersensitivity to BCG purified proteins derivatives as shown by a

negative skin-test. The mice developed a small sensitization against PPD only after injection of large dose of extended freeze-dried killed BCG (1 mg and 10 mg). By comparison, mice immunised with living BCG or heat-killed BCG, even at low doses (10^7 CFU of living BCG or equivalent to 10^8 bacterial corpses for heat-inactivated BCG) develop a delayed type hypersensitivity to BCG purified proteins derivatives as shown by a positive skin-test. These results are correlated with the preceding results (figures 9 and 21) showing a small production of IFN- γ under specific stimulation, in the extended freeze-dried killed BCG treated groups, contrary to the groups treated with other BCG preparations showing production of higher level of IFN- γ under specific stimulation. Similarly, reactivity of guinea-pigs to PPD is marginal after extended freeze-dried killed BCG treatment.

These results indicate that following immunisation with extended freeze-dried killed BCG, a particular sub-type of specific T lymphocytes are selected or alternatively specific T cells are immobilised in some particular tissues. Consequently, as opposed to immunisation with living BCG or heat-killed BCG, immunisation with extended freeze-dried killed BCG does not interfere with the diagnosis of tuberculosis by the DTH skin-test.

EXAMPLE 6: PROTECTIVE EFFECT OF EXTENDED FREEZE-DRIED (EFD) KILLED MYCOBACTERIA ON ASTHMA IN A GUINEA-PIG MODEL.

1) Material and methods

Male adult guinea-pigs (350 g) were immunized by the subcutaneous route with 10 μ g ovalbumin (OVA; ICN Laboratories) and 1 mg alum in saline (0.4 ml final volume).

Three weeks after, said guinea-pigs (3 groups of 10 animals) were injected intradermally with either extended freeze-dried killed BCG (10^7 or 10^8 CFU equivalent *i.e.* 10 μ g or 100 μ g), or PBS (control).

The broncho-pulmonary reactivity is assayed on guinea-pigs with a method different to that used for mice; the sensitivity of each animal was evaluated with increasing doses of aerosolized histamine (20, 50, 100 or 400 μ g), 24 h before

allergen challenge, and the highest concentration of histamine giving elevated Penh values was considered to be the basal level of sensitivity to histamine.

A similar measure was performed 24 h after the allergen challenge; OVA immunized guinea pigs develop a higher sensitivity to histamine, less histamine
5 is thus needed to obtain the same Penh values than before the challenge.

More precisely:

Six guinea-pigs of each group have been tested for their broncho-pulmonary reactivity with an histamine test performed 7 to 10 weeks after the injection of BCG or control product (e.g. 10 to 13 weeks after the ovalbumin immunisation).
10

The dosage of histamine triggering a clear bronchoconstriction is measured, for each guinea-pig, with a barometric plethysmograph.

Each guinea-pig is its own control, knowing that the basal reactivity is determined the 1st day of the test.

15 24 hours after, an ovalbumin aerosol is administered. The 3rd day the histamine reactivity is measured again.

The administration of ovalbumin increases the broncho-pulmonary hyper-reactivity to histamine.

2) Results

20 The results presented in figure 14 show that 5/6 and 6/6 guinea-pigs that have been treated respectively with 10 µg (10⁷ CFU equivalent) (figure 14B) or 100 µg (10⁸ CFU equivalent) (figure 14C) extended freeze-dried killed BCG according to the invention, are protected against broncho-pulmonary hyper-reactivity (no increase of the histamine sensitization) whereas in the control group (figure 14A)
25 (no treatment with BCG) there is a significant increase of broncho-pulmonary hyper-reactivity to histamine (x4).

EXAMPLE 7: MINIMAL SIDE EFFECTS OF EXTENDED FREEZE-DRIED KILLED BCG

a) No anemia and no thrombopenia

30 Mice were injected intravenously with extended freeze-dried killed BCG (10 µg) or the different BCG preparations as described in example 3d, or non-

treated. 18 days after the injection, red blood cells and platelets numbers were determined from a blood sample.

The results presented in figure 22 show that, contrary to the living BCG and the heat-killed BCG treated groups, no anemia and no thrombopenia were observed in the extended freeze-dried BCG treated group.

b) Minimal inflammatory reaction at the site of injection

At day 0, the different BCG preparations, as described in example 3d, were injected subcutaneously (in the footpad area) to groups of mice. Living BCG was injected once only and heat-killed and extended freeze-dried killed BCG were injected twice (day 0 and day 36). The foot pad increase was then measured every week for ten weeks.

The results presented in figure 23 show that extended freeze-dried killed BCG exhibits minimal inflammatory side effects.

c) Reduced TNF- α production after LPS stimulation

Lung explants from mice treated 18 days previously with the different BCG preparations as described above, were maintained *in vitro* in presence or not of LPS, and TNF- α production was measured by ELISA using commercial kits, according to the manufacturer's instructions. The results presented in figure 24 show that, contrary to living BCG and heat-killed BCG treatment, extended freeze-dried killed BCG treatment does not induce the production and/or the activation of macrophages.

EXAMPLE 8 : ANALYSIS OF THE CELLS IN THE DRAINING LYMPH NODES AND IN THE SPLEEN AFTER SUBCUTANEOUS INJECTION OF EXTENDED FREEZE-DRIED KILLED BCG.

1) **Material and Methods**

The cells present in the draining lymph nodes of mice treated 48 hours previously by the subcutaneous route, with the different BCG preparations as described in example 3d, were analysed by flow cytometry using monoclonal antibodies directed to cell surface markers of different leukocyte populations, labelled with various fluorochromes.

The cells present in the spleen of mice treated by the subcutaneous route with extended freeze-dried killed BCG, were analysed as above described, after *in vitro* stimulation in the presence of OVA or BCG culture supernatant or no stimulation.

5 2) Results

a) draining lymph nodes

The analysis of the cells present in the draining lymph nodes show that by comparison with the living BCG and the heat-killed BCG treated groups, the cells present in the lymph nodes were three to five times more numerous 48 h after subcutaneous injection of extended freeze-dried killed BCG. A kinetic analysis of the different leukocyte populations for four days after the injection of extended freeze-dried killed BCG, show a transient increase of the dendritic cells (CD11c⁺) number at 48 hours, as well as an increasing number of B220⁺ and CD4⁺ lymphocytes from 6 hours to 96 hours.

15 b) spleen

The analysis of the cells present in the spleen show:

- a significant increase in the number of CD11c⁺ Gr1⁺ B220⁺ plasmacytoid dendritic cells in the extended freeze dried treated group, no increase of the number of CD8α⁺ cells was observed in the same group (figure 25).
- 20 - a significant increase in the number of CD4⁺ CD25⁺ IL-10⁺ cells in the extended freeze dried treated group, either stimulated *in vitro* with OVA or BCG culture supernatant or non-stimulated (figure 26).

The results show that in the mice model of asthma, extended freeze-dried killed BCG is able to induce a significant increase in the number of cells which induce immunomodulatory cells (CD11c⁺ Gr1⁺ B220⁺ plasmacytoid dendritic cells) or are themselves immunomodulatory cells (CD4⁺ CD25⁺ IL-10⁺ cells) in the mice which have been previously sensitized to an allergen (OVA or water soluble ray grass pollen), which results in a protective effect against the symptoms of asthma.

EXAMPLE 9 : Extended freeze-dried killed BCG is active by the oral route

30 Mice which have been immunized previously with OVA, as described in example 3d were treated orally with 1 mg extended freeze-dried at days

42, 44, 46 and 62, 64 and 66 or non-treated and then challenged or not with OVA at day 96. The protective effect of extended freeze-dried killed BCG was assessed by numeration of lungs cells infiltrate as described in example 3.

5 The results show a significant decrease $p < 0.001$ of the total cell number in the lungs of the extended freeze-dried killed BCG treated group, by comparison with the untreated group (figure 27). Analysis of the leukocyte populations show a significant decrease of the macrophages, polynuclears and dendritic cells numbers in the lungs of the treated group.

EXAMPLE 10 : Determination of the active dose, the rhythm of administration,
10 **the duration and the delay of action of extended freeze-dried killed BCG**

Extended freeze-dried killed BCG was injected by the subcutaneous route to mice previously immunized and then challenged with OVA, and its activity was then tested as described in example 3.

a) Active dose and dose number

15 Adult mice were sensitized with OVA at days 0 and 7, injected by the subcutaneous route with extended freeze-dried killed BCG once at day 45 or twice at days 45 and 65, and then challenged with ovalbumine at day 96. The activity of extended freeze-dried killed BCG was tested by the prevention of bronchopulmonary hyper-reactivity in the BP2 model of asthma as described in example 3.

20 The smallest dose (10 μg) was active and doses larger than 100 μg did not increase the activity.

A protective effect was observed with one dose (10 μg) of extended freeze-dried killed BCG only (figure 28).

b) Delay of action of extended freeze-dried killed BCG

25 In order to determine the time sequence to obtain the extended freeze-dried killed BCG efficacy, adult mice (6 per group) were sensitized with OVA at days 0 and 7, injected subcutaneously with 100 μg extended freeze-dried killed BCG at day 14 and then challenged with OVA at day 21, 28 or 35. The activity of extended freeze-dried killed BCG was tested by the prevention of bronchopulmonary
30 hyper-reactivity in the BP2 model of asthma as described in example 3.

The prevention of bronchopulmonary hyper-reactivity occurs only if a delay of two to three weeks is present between extended freeze-dried killed BCG administration and challenge (figure 29) ; this result indicates that the recruitment and expansion of a given cell population is required before extended freeze-dried killed BCG is active. This result confirms also that one dose of extended freeze-dried killed BCG is sufficient to protect the mice against asthma.

c) Duration of action of extended freeze-dried killed BCG

In order to determine the duration of action of extended freeze-dried killed BCG, adult mice (6 per group) were sensitized with OVA at days 0 and 7, injected subcutaneously with 100 µg extended freeze-dried killed BCG at day 45 and 65 and then challenged with OVA at day 96 or 42. The activity of extended freeze-dried killed BCG was assessed by the preventive effect on bronchopulmonary hyper-reactivity and by numeration of the lungs cell infiltrate, in the BP2 model of asthma, as described in example 3.

The results show that the extended freeze-dried killed BCG protective effect persists at least for two months (figure 30).

d) preventive versus curative protocol of administration of extended freeze-dried BCG.

The previous results as presented above, demonstrate the efficacy of extended freeze-dried killed BCG in a "curative protocol" (administration after allergen immunization). Thus, in order to evaluate the efficacy of extended freeze-dried killed BCG in a preventive protocol, mice were treated as follows:

- adult mice (6 to 7 weeks old; 6 mice per group) were injected subcutaneously with extended freeze-dried killed BCG (100 µg) at days 14, 21 or 28 before OVA immunization (days-14, -21 and -28), immunized with OVA at days 0 and 7 and then challenged with OVA at day 14.

- newborn mice (7 to 8 days old, 6 to 8 mice per group) were injected subcutaneously or intranasally with extended freeze-dried killed BCG (100 µg) at day 56 before OVA immunization (day -56), immunized with OVA at days 0 and 7 and then challenged with OVA at day 14.

The activity of extended freeze-dried killed BCG in the preventive protocol was assessed by the preventive effect on broncho-pulmonary hyper-reactivity in the BP2 model of asthma, as described in example 3.

5 The results in adult mice show a protective effect of extended freeze-dried BCG in a preventive protocol, only when the preparation is injected some days before sensitization (2 weeks and 3 weeks).

The results in newborn mice show a protective effect of extended freeze-dried BCG in a preventive protocol, only with the subcutaneous administration; the intranasal route is poorly active in these conditions.

CLAIMS

1°) A bacterial preparation, characterized in that:

- it contains killed Gram positive bacteria, obtainable by a process which does not denature the structure of the molecules from the bacteria cells, and

5 - it is able to induce, *in vivo*, a modulation of the immune response against an antigen.

2°) The bacterial preparation according to claim 1, characterized in that said Gram positive bacteria are chosen from Gram positive facultative intracellular bacteria.

10 3°) The bacterial preparation according to claim 2, characterized in that said Gram positive facultative intracellular bacteria are selected from the group consisting of *Listeria sp.*, *Corynebacterium sp.*, and Actinomycetes comprising *Mycobacteria sp.*, *Nocardia sp.* and *Rhodococcus sp.*.

15 4°) The bacterial preparation according to claim 3, characterized in that said mycobacteria is *Mycobacteria bovis*, more preferably *Mycobacteria bovis* BCG.

5°) The bacterial preparation according to anyone of claims 1 to 4, characterized in that it contains extended-freeze-dried killed bacteria obtainable by an extended freeze-drying process.

20 6°) The bacterial preparation according to claim 4, characterized in that it contains extended freeze-dried killed mycobacteria.

7°) The bacterial preparation according to claim 5 or claim 6, characterized in that it contains less than 1.5% of residual water, preferably less than 1 %, more preferably less than 0.5%.

25 8°) The bacterial preparation according to anyone of claims 5 to 7, characterised in that it is prepared by an extended freeze-drying process, comprising at least the steps of :

(i) harvesting a culture of live bacteria cells,

(ii) washing the bacteria cells in water or in an aqueous solution of a salt such as borate,

30 (iii) freezing the bacteria cells in water or in an aqueous solution of a salt such as borate,

(iv) killing the frozen bacteria cells by drying them in a lyophiliser, for a time sufficient to remove at least 98.5 % of the water, preferably at least 99 % of the water, more preferably at least 99.5 % of the water, and

(v) collecting the extended freeze-dried killed bacteria cells.

5 9°) The bacterial preparation according to anyone of claims 5 to 7, characterized in that it is prepared by an extended freeze-drying process comprising the steps (i), (iii), (iv) and (v) as defined in claim 8.

10 10°) The bacterial preparation according to claims 8 or 9, characterized in that it is prepared by performing step (iv) at a drying chamber pressure of about 0.02 mBar to 0.2 mBar; more preferably of about 0.06 mBar to 0.1 mBar.

 11°) Fractions of bacterial preparation according to claims 1 to 10, selected in the group consisting of:

- a fraction A consisting of an organic solvent extract of said killed bacterial preparation,
- 15 - a fraction B consisting of a glycosidase-treated extract of said killed bacterial preparation,
- a fraction C consisting of a DNase and/or a RNase-digested extract of said killed bacterial preparation,
- a fraction D, consisting of a protease-treated extract of said killed bacterial preparation,
- 20 - a fraction E, consisting of a extract of said killed bacterial preparation successively treated by an organic solvent, a glycosidase, a DNase and/or a RNase, and finally a protease.

 12°) Fractions according to claim 11, characterized in that:

25 - said fraction A is obtained by extraction with a methanol/chloroform mixture,

 - said fraction B is obtained by digestion with lysozyme,

 - said fraction C is obtained by digestion with DNase I and/or RNase A,

 and

30 - said fraction D is obtained by digestion with subtilisine.

13°) A pharmaceutical composition, characterized in that it comprises an effective amount of a bacterial preparation according to anyone of claims 1 to 10 and/or at least one fraction thereof, according to claim 11 or claim 12, a pharmaceutically acceptable carrier and/or an additive, and/or an immunostimulant, and/or an adjuvant and/or an immunomodulator distinct from the bacterial preparation according to claim 1.

14°) Use of the bacterial preparation according to anyone of claims 1 to 10 or fraction(s) thereof according to claims 11 or 12 for the preparation of a medicament for the prevention and/or the treatment of diseases comprising an immune dysregulation.

15°) Use according to claim 14, wherein said diseases comprising an immune dysregulation are selected from the group consisting of cancer, autoimmune diseases and allergy.

16°) Use according to claim 15, wherein said autoimmune diseases are selected from the group consisting of multiple sclerosis, rheumatoid arthritis, Crohn disease and diabetes mellitus,

17°) Use according to claim 15, wherein said allergy is selected from the group consisting of asthma, allergic rhinitis, conjunctivitis and atopic dermatitis.

18°) Use according to anyone of claims 14 to 17, wherein said medicament is intended to be administered orally, sublingually, parenterally and preferably subcutaneously or intranasally.

19°) A method for the preparation of the extended freeze-dried killed bacterial preparation as defined in anyone of claims 5 to 10, characterized in that it comprises at least the steps of:

- (i) harvesting a culture of live bacteria cells,
- (ii) washing the bacteria cells in water or in an aqueous solution of a salt,
- (iii) freezing the bacteria cells in water or in an aqueous solution of a salt,
- (iv) killing the frozen bacteria cells by drying them in a lyophiliser, for a time sufficient to remove at least 98.5% of the water, preferably at least 99 % of the water, more preferably at least 99.5 % of the water, and
- (v) collecting the extended freeze-dried killed bacteria cells.

20°) A method for the preparation of the extended freeze-dried killed bacterial preparation as defined in anyone of claims 5 to 10, characterized in that it comprises the steps (i), (iii), (iv) and (v) as defined in claim 19.

21°) The method of claims 19 or 20, characterised in that step (iv) is performed at a drying chamber pressure of about 0.02 mBar to 0.2 mBar, more preferably 0.06 mBar to 0.1 mBar.

22°) Kit for the treatment of a disease comprising an immune dysregulation, characterized in that it comprises at least a pharmaceutical composition according to claim 13, means of administration of said pharmaceutical composition and/or means for insuring compliance with the prescribed treatment.


23°) Products containing a bacterial preparation according to anyone of claims 1 to 10 or fractions thereof according to claim 11 or claim 12 and a product selected from the group consisting of anticancer, anti-diabetes and immunomodulatory drugs as a combined preparation for simultaneous, separate or sequential use in the prevention and/or the treatment of diseases comprising an immune dysregulation.

24°) Products according to claim 23, characterized in that said product is an anti-histaminic drug or an anti-inflammatory drug.


25°) Use of a bacterial preparation according to anyone of claims 1 to 10 or fractions thereof according to claim 11 or claim 12 for the preparation of a medicament for the treatment of asthma to be administered by the oral, parenteral, sublingual or intranasal route, in a dose range which is equivalent in mice to 1 µg to 10 000 µg, preferably 10 µg to 1 000 µg, most preferably 10 µg to 100 µg, at two months intervals, starting from at least two weeks before the usual period of allergen exposure.

26°) Use of a bacterial preparation according to anyone of claims 1 to 10 or fractions thereof according to claim 11 or claim 12 for the preparation of a medicament for the prevention of asthma in human baby, to be administered by the oral, parenteral, sublingual or intranasal route, in a dose range which is equivalent in mice to 1 µg to 10 000 µg, preferably 10 µg to 1 000 µg, most preferably 10 µg to 100 µg.

1/34



Killing by freeze drying



Vaccine production

FIGURE 1

2/34




FIGURE 2

3/34




FIGURE 3

4/34




FIGURE 4

5/34




FIGURE 5

6/34




FIGURE 6

7/34




FIGURE 7

8/34






FIGURE 8



10/34



IgE specific against OVA




FIGURE 10

12/34




FIGURE 11A

13/34




FIGURE 11B

14/34




FIGURE 12A

15/34




FIGURE 12B

16/34

DTH toward PPD




FIGURE 13

17/34




FIGURE 14B




FIGURE 14A

18/34




FIGURE 14C




Figure 15

20/34

A

B





Figure 16

21/34




Figure 17

22/34




Figure 18

23/34





Figure 19



25/34




Figure 21




Figure 22

27/34




Figure 23

28/34




Figure 24




Figure 25

30/34




Figure 26

31/34




Figure 27




Figure 28




Figure 29




Figure 30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 June 2003 (19.06.2003)

PCT

(10) International Publication Number
WO 03/049752 A3

(51) International Patent Classification⁷: **A61K 35/74**,
39/04, A61P 35/00, 37/00, 31/06

(21) International Application Number: **PCT/IB02/05760**

(22) International Filing Date:
11 December 2002 (11.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01403195.9 11 December 2001 (11.12.2001) EP

(71) Applicants (for all designated States except US): **INSTITUT PASTEUR** [FR/FR]; 28 rue du Docteur Roux, F-75724 Cedex 15 Paris (FR). **INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE** [FR/FR]; 101 rue de Tolbiac, F-75654 Cedex 13 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NAHORI, Marie-Anne** [FR/FR]; 147 avenue Henri Ginoux, F-92120 Montrouge (FR). **LAGRANDERIE, Micheline** [FR/FR]; 84 rue Charles Laffitte, F-92200 Neuilly-sur-Seine (FR). **MARCHAL, Gilles** [FR/FR]; 4 rue Francisco Ferrer, F-94200 Ivry-sur-Seine (FR). **VARGAFTIG, Bernardo, Boris** [FR/FR]; 5 Villa Poirier, F-75015 Paris (FR). **LEFORT, Jean** [FR/FR]; 2 rue des Cailloux, F-92110 Clichy-sur-Seine (FR). **ROMAIN, Félix** [FR/FR]; 49 bis rue Dreyfus, F-91640 Fontenay les Briis (FR).

HEKIMIAN, Georges [FR/FR]; 91 rue Jean Mermoz, F-92380 Garches (FR). **PELTRE, Philippe** [FR/FR]; 100 Boulevard Georges Seurat, F-92200 Neuilly-sur-Seine (FR).

(74) Agent: **CABINET ORES**; 36 rue de St Petersburg, F-75008 Paris (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
20 November 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GRAM POSITIVE BACTERIA PREPARATIONS FOR THE TREATMENT OF DISEASES COMPRISING AN IMMUNE DYSREGULATION

(57) Abstract: Compositions comprising components prepared from Gram positive bacteria such as Gram positive facultative intracellular bacteria, for example mycobacteria, including extended freeze-dried killed Gram positive bacteria, their method of preparation and their use in humans and animals, for the prevention and/or the treatment of disorders comprising an immune dysregulation such as cancer, autoimmune diseases, allergy and tuberculosis.

WO 03/049752 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 02/05760

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K35/74 A61K39/04 A61P35/00 A61P37/00 A61P31/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE, BIOSIS, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 1 027 979 A (CIBA LTD) 4 May 1966 (1966-05-04) page 2, line 34 - line 45; example 4 ---	1-10, 13, 14, 19-22
X	GB 764 718 A (GLAXO LAB LTD) 2 January 1957 (1957-01-02) page 3, line 13 - line 15 ---	1-10, 13, 14, 19-22
X	US 3 594 471 A (HERTZBERGER ELLIS ET AL) 20 July 1971 (1971-07-20) the whole document column 3, line 15 - line 23; claims --- -/--	1-6

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

5 August 2003

Date of mailing of the international search report

26/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Bayrak, S

INTERNATIONAL SEARCH REPORT

In national Application No
PCT/IB 02/05760

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KYD J M ET AL: "Killed whole bacterial cells, a mucosal delivery system for the induction of immunity in the respiratory tract and middle ear: an overview" VACCINE, BUTTERWORTH SCIENTIFIC. GUILDFORD, GB, vol. 17, no. 13-14, January 1999 (1999-01), pages 1775-1781, XP004158321 ISSN: 0264-410X the whole document	1-26
A	ALEXANDROFF A B ET AL: "BCG immunotherapy of bladder cancer: 20 years on" LANCET, XX, XX, vol. 353, no. 9165, 15 May 1999 (1999-05-15), pages 1689-1694, XP004266316 ISSN: 0140-6736 the whole document	1-26

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 02/05760

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: claims 14-18 (all partially)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: claims 14-18 (all partially)

Claims 14-18 relate to the use freeze-dried killed mycobacteria preparation for use in the treatment of diseases comprising an "immune dysregulation" and "cancer, autoimmune diseases and allergy", which encompass a multitude of different diseases. The claims thus cover a rather large number of diseases, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of diseases. In the present case, the, the claims so lack support and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims 2-17 also lack clarity because it is not fully possible to determine the diseases for which protection might legitimately be sought (Article 84 EPC). Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely asthma, tuberculosis, multiple sclerosis, rheumatoid arthritis, Crohn disease, diabetes mellitus, allergisch rhinitis, conjunctivitis, and atopic dermatitis.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/IB 02/05760

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1027979	A	04-05-1966	CH 465768 A	30-11-1968
			AT 257050 B	25-09-1967
			BE 650404 A	11-01-1965
			DE 1442075 A1	21-11-1968
			DK 106687 C	06-03-1967
			FR 1419643 A	03-12-1965
			NL 6407893 A	13-01-1965
			SE 333432 B	15-03-1971
			US 3567585 A	02-03-1971
			AT 253121 B	28-03-1967
GB 764718	A	02-01-1957	NONE	
US 3594471	A	20-07-1971	AT 282824 B	10-07-1970
			BE 723727 A	12-05-1969
			DE 1807820 A1	07-08-1969
			DK 121454 B	18-10-1971
			ES 360080 A1	01-10-1970
			FR 8174 M	31-08-1970
			GB 1163470 A	04-09-1969
			NL 6816071 A	16-05-1969

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.